

# ON THE LOCAL STRUCTURE OF UNIQUELY 3-COLORABLE PLANE GRAPHS

TOMÁŠ MADARAS

*Pavol Jozef Šafárik University in Košice*

e-mail: tomas.madaras@upjs.sk

ALFRÉD ONDERKO, ROMAN SOTÁK AND DIANA ŠVECOVÁ

*Pavol Jozef Šafárik University in Košice*

e-mail: alfred.onderko@upjs.sk, roman.sotak@upjs.sk,

diana.svecova@student.upjs.sk

A graph  $G$  is called uniquely  $k$ -colorable if  $\chi(G) = k$  and every  $k$ -coloring of  $G$  induces the same partition of the vertex set of  $G$  into  $k$  independent sets. It is known that uniquely 3-colorable plane graphs possess interesting structural properties: except of  $C_3$ , they contain at least three triangles ([1]) and in the case they contain exactly four triangles, two of them are adjacent ([3]). Moreover, they contain a 3-face adjacent to a face of size at most 5 ([2]).

In addition to these results, we show that every uniquely 3-colorable plane graph of minimum degree at least 3 contains an edge of type  $(3, \leq 9)$ ,  $(4, \leq 6)$  or  $(5, 5)$  (the bounds being sharp); a similar result holds for 3-vertex paths (with all vertices having degrees at most 13), but not for longer paths. Also, extending the result of [2], we prove that every uniquely 3-colorable plane graph contains a cluster of three faces of sizes at most 7 as well as a cluster of four faces of size at most 9 (this bound being sharp).

## References

- [1] V.A. Aksionov, On uniquely 3-colorable planar graphs, *Discrete Mathematics* 20 (1977), 209–216.
- [2] Z. Li, N. Matsumoto, E. Zhu, J. Xu, T. Jensen, On uniquely 3-colorable plane graphs without adjacent faces of prescribed degrees, *Mathematics* 7(9) (2019) 793
- [3] Z. Li, E. Zhu, Z. Shao, J. Xu, A note on uniquely 3-colourable planar graphs, *International Journal of Computer Mathematics* 94(5) (2017), 1028–1035.